

Perturbation enstrophy decay in Poiseuille and Couette flows according to Synge's method

Original

Perturbation enstrophy decay in Poiseuille and Couette flows according to Synge's method / Domenicale, Loris; Fraternale, Federico; Staffilani, Gigliola; Tordella, Daniela. - In: BULLETIN OF THE AMERICAN PHYSICAL SOCIETY. - ISSN 0003-0503. - STAMPA. - 68:(2015), pp. 249-249. (Intervento presentato al convegno 68th American Physical Society - Division of Fluid Dynamics Annual Meeting 2015 tenutosi a Boston nel 22-24 Novembre 2015).

Availability:

This version is available at: 11583/2657307 since: 2017-09-22T13:07:10Z

Publisher:

American Physical Society

Published

DOI:

Terms of use:

openAccess

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

(Article begins on next page)

Bulletin of the American Physical Society

68th Annual Meeting of the APS Division of Fluid Dynamics Volume 60, Number 21

Sunday–Tuesday, November 22–24, 2015; Boston, Massachusetts

Session G16: Flow Instability: General II

8:00 AM–10:10 AM, Monday, November 23, 2015

Room: 204

Chair: Nathaniel Barlow, Rochester Institute of Technology

Abstract ID: BAPS.2015.DFD.G16.2

Abstract: G16.00002 : Perturbation Enstrophy Decay in Poiseuille and Couette Flows according to Synge's Method

8:13 AM–8:26 AM

[Preview Abstract](#)

[MathJax On](#) | [Off](#) [← Abstract](#) [→](#)

Authors:

Loris Domenicale

(Politecnico di Torino, Matematica per le Scienze dell'Ingegneria)

Federico Fraternali

(Politecnico di Torino, DIMEAS)

Gigliola Staffilani

(MIT, Department of Mathematics)

Daniela Tordella

(Politecnico di Torino, DIMEAS)

In this work we derive the conditions for no enstrophy growth for bidimensional perturbations in the plane Couette and Poiseuille flows. We follow the method of vorticity proposed by Synge in 1938 (see the Semi-Centennial Publication of the Amer. Math. Soc., equation 12.13, and the more detailed version in the Proc. of the Fifth Inter. Congress of Applied Mechanics, pages 326-332), which is actually based on the analysis of the spatially averaged enstrophy. We find that the limit curve in the perturbation wavenumber-Reynolds number map differs from the limit for no energy growth (see e.g. Reddy 1993). In particular, the absolute stability region for the enstrophy is wider than that of the kinetic energy, and the maximum Reynolds number giving the monotonic enstrophy decay, at all wavenumbers, is 155 and 80 for the Poiseuille and Couette flows, respectively. It should be noted that in past literature the energy-based analysis was preferred to Synge's enstrophy analysis. This, possibly, for two reasons: the low diffusivity of the 1938 Vth ICAM proceedings and the objectively very complicated analytical treatment required. Nevertheless, the potentiality of this method seems high and therefore it is interesting nowadays to exploit it by means of the symbolic calculus.

To cite this abstract, use the following reference: <http://meetings.aps.org/link/BAPS.2015.DFD.G16.2>